

Sample Standard Deviation

$$S = \sqrt{\frac{\sum (x - \bar{x})^2}{n - 1}}$$

Population Standard Deviation

$$\sigma = \sqrt{\frac{\sum (x - \mu)^2}{N}}$$

Sample Standard Deviation without calculating the mean

$$s = \sqrt{\frac{n (\sum x^2) - (\sum x)^2}{n (n - 1)}}$$

Mean of grouped Data

$$\bar{x} = \frac{\sum f \cdot x}{n}$$

NOTE

Variance = $(\sigma)^2$
OR $(s)^2$

Standard Deviation of frequency table or grouped data


$$S = \sqrt{\frac{n [\sum (f \cdot x^2)] - [\sum (f \cdot x)]^2}{n (n - 1)}}$$

Standard Deviation of frequency table or grouped data where x_m is the midpoint of the class interval

$$S = \sqrt{\frac{n [\sum (f \cdot x_m^2)] - [\sum (f \cdot x_m)]^2}{n (n - 1)}}$$


Skewness

Figure 2-13 (b)




Mode = Mean = Median

SYMMETRIC




Mean Median Mode

SKEWED LEFT (negatively)



Mode Median Mean

SKEWED RIGHT (positively)



$$\text{Percentile score} = \frac{\text{number of scores below data value} + 0.5}{\text{total number of data values}} \times 100$$

Finding a Data Value Corresponding to a given Percentile

$$c = \frac{n \cdot p}{100}$$

where c = data value position

n = total number of values

p = percentile

-if c is a whole number count go to the number between c and $c+1$

-if c is not a whole number round up

Outliers
 Interquartile Range (or IQR): $Q_3 - Q_1$
 To Check for Outliers: $Q_3 + 1.5 \text{ IQR}$
 $Q_1 - 1.5 \text{ IQR}$

BOXPLOT
5 - number summary

- Minimum
- first quartile Q_1
- Median (Q_2)
- third quartile Q_3
- Maximum

RANGE: Highest - Lowest
MIDRANGE: $\frac{\text{Highest} + \text{Lowest}}{2}$

Probability Distribution

mean of a probability distribution $\mu = \Sigma [x \cdot P(x)]$

variance of probability distribution $\sigma^2 = [\Sigma x^2 \cdot P(x)] - \mu^2$

standard deviation of probability distribution $\sigma = \sqrt{[\Sigma x^2 \cdot P(x)] - \mu^2}$

Expected Value $E = \mu = \Sigma [x \cdot P(x)]$
 The average value of outcomes

Binomial Distribution

Mean of a binomial distribution $\mu = n \cdot p$

Standard deviation of a binomial distribution $\sigma = \sqrt{n \cdot p \cdot q}$

Binomial Probability Formula $P(X) = {}_n C_x \cdot p^x \cdot q^{n-x}$

Central Limit Theorem

the mean of the sampling distribution $\mu_{\bar{x}} = \mu$

the standard deviation of the sampling distribution (standard error) $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$

central limit theorem formula for z-score $Z = \frac{\bar{X} - \bar{\mu}}{\sigma / \sqrt{n}}$

Population z-score
 $Z = \frac{x - \mu}{\sigma}$

To find the x-value:
 $x = z\sigma + \mu$

Confidence Interval

$$\bar{X} \pm Z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}$$

Confidence Interval for t-distribution

$$\bar{x} \pm t_{\alpha/2} \frac{s}{\sqrt{n}}$$

Confidence Interval for Proportion

$$\hat{p} \pm z_{\alpha/2} \sqrt{\frac{\hat{p}\hat{q}}{n}}$$

Margin of error (E)
or Error Estimate:

$$Z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}$$

Sample size

$$n = \left(\frac{Z_{\alpha/2} \sigma}{E} \right)^2 \quad \text{or} \quad n = \frac{(Z_{\alpha/2})^2 \hat{p} \hat{q}}{E^2}$$

Note: Point estimate is the mean

Hypothesis Testing: null always contains the equality
p-value method and critical value method

Test Statistic

$$z = \frac{\bar{x} - \bar{\mu}}{\sigma / \sqrt{n}}$$

Test Statistic for a proportion

$$Z = \frac{\hat{p} - p}{\sqrt{\frac{pq}{n}}}$$

Contingency Tables Chi Square test of Independence

Expected (E): $\frac{(\text{row total}) \times (\text{column total})}{\text{grand total}}$

Test of Independence

Test Statistic

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

H₀: row variable is independent of column variable

H_a: row variable is dependent on column variable

Critical Values

- Found in Table G using **degrees of freedom = (r - 1)(c - 1)**
r is the number of rows
c is the number of columns
- Tests of Independence are always right-tailed.

	H ₀ True Innocent	H ₀ False Not Innocent Guilty
Reject H ₀ Convict	Type 1 Error	correct decision
Fail to Reject H ₀ Not Convict	correct decision	Type 2 Error